



## **Evaluation of the diversity and the parasitic load of the waters of the Bouishak Wadi in the city of Meknes (Central Morocco)**

**K. Ouarrak<sup>1</sup>, A. Chahlaoui<sup>1</sup>, H. El Omari<sup>1</sup>, I. Taha<sup>1</sup>, A. Taouraout<sup>1,2</sup>, A. Kharroubi<sup>3</sup>**

<sup>1</sup> Equipe Gestion et valorisation des ressources naturelles, Laboratoire Santé et environnement, Faculté des Sciences, Université Moulay Ismail, Meknès, Maroc

<sup>2</sup> Laboratory of Agrophysiology, Biotechnology, Environment and Qualities, Faculty of Sciences, University Ibn Tofail of Kenitra, P.O. Box 14000 Morocco.

<sup>3</sup> RU: Applied Hydrosciences Research Unit, Higher Institute of Water Sciences and techniques, University of Gabès, P.O. Box 6072 Tunisie

Received 13 July 2019,  
Revised 10 Oct 2019,  
Accepted 15 Oct 2019

### **Keywords**

- ✓ Wastewater
- ✓ Bouishak Wadi
- ✓ Parasites,
- ✓ Meknes ,
- ✓ Morocco.

[kouarrak@yahoo.com](mailto:kouarrak@yahoo.com) ;  
Phone: +21245366633

### **Abstract**

The objective of this study is to evaluate the diversity and abundance of the forms of dissemination of helminths and protozoa present in the waters of wadi Bouishak (Meknes, Morocco), and to elucidate the risks that these parasites present for health. The wastewater sampling has carried out monthly during a period from January to December 2017 between 10 am and 11 am at the three stations (B1, B2 and B3) that flow into the wadi. The 36 samples analyzed were positive with a predominance of protozoan cysts on Helminth eggs ( $29.08 \cdot 10^5$  cysts/L against 4573 eggs/L). Thirteen parasite taxa has detected: *Giardia lamblia*, *Entamoeba histolytica*, *Entamoeba coli*, *Blastocystis hominis*, *Ascaris lumbricoides*, *Trichuris trichiura*, *Ancylostoma duodenale*, *Capillaria sp* *Toxocara canis*, *Enterobius vermicularis*, *Taenia saginata*, *Hymenolepis nana* and *Hymenolepis diminuta*. The results obtained reveal a high parasite contamination of the watercourse, which could have negative effects on the quality of the environment in general and on human and animal health in particular.

## **1. Introduction**

Demographic, economic and climatic constraints lead to an increase in the demand for water, which causes a great pressure on the surface water. Yet the surface waters has also polluted by industrial, domestic and agricultural waste. In Morocco wastewater is used as an important resource for agriculture that suffers from water deficit [1-2]. Without prior control, these effluents pose such a significant threat to the environment, human and animal health [3-4]. According to WHO, serious health problems has caused by waterborne diseases [5]. In the region of Meknes, wastewater has discharged without prior treatment into the watercourses crossing the city (Ouislane wadi, Boufakrane wadi and Bouishak wadi). As a result, these rivers has considered as open sewers through urban and peri-urban areas of the city. The reuse of these raw or mixed effluents in irrigation is a real practice. Previous studies have examined the impact of liquid effluents on the water quality of these rivers [6-7-8].


The choice of the Bouishak valley lies in the fact that it constitutes an extreme case since the raw sewage represents 93% of the water used for irrigation [9]. The bed of the Bouishak wadi drains wastewater from densely populated areas (Hay El Mansour, Kamilia and Toulal). Agriculture is mainly oriented towards market gardening and arboriculture. In the current state of knowledge, no parasitological study has carried out. The objective of this study is to evaluate the diversity and abundance of the forms of dissemination of Helminths and Protozoa that are

present in the waters of the Bouishak wadi and to elucidate the health risk emanating from these parasites. To do this, spatio-temporal monitoring of the parasitological quality of sewage discharged into the Bouishak wadi has carried out.

## 2. Material and Methods

### 2.1. Study area

The city of Meknes is located in the northern part of Morocco. It occupies an area of about 79210 km<sup>2</sup>. The study area belongs to the Fez-Meknes region according to the new 2015 administrative division (Figure 1). It is the second city of the plain of Sais, after the city of Fez. This area has characterized by geological diversity consisting of hard lake limestones and tawny sands. The climate is Mediterranean, with an average rainfall of 660 mm / year [10]. According to the 2014 General Census of Population and Housing [11], the population of the city of Meknes is in the order of 835,695 inhabitants against 715,284 in 2004. As a result, the average annual growth rate reached 1.6% for the period 2004-2014.




**Figure 1:** Geographical location of the study area

### 2.2. Sampling and analysis methods

For a more representative sampling of the watercourse, we have chosen three stations on the Bouishak Wadi bed (Figure 2): Station B1 is located upstream of watercourses in a zone of great anthropogenic influence (domestic, industrial and agricultural), stations B2 and B3 are located downstream of the agglomeration of Meknes (Table 1). Monthly samples were taken between 10 am and 11 am, when the parasite load is closely related to human activity [12-13- 14], during a period from January to December 2017. À chaque station, un échantillon de 1 litre d'eaux usées a été prélevé et conservé en ajoutant du formol, 10% (2 ml / 1 d'eaux usées) étiqueté et transporté dans un réfrigérateur (à une température de + 4 ° C) au laboratoire d'hydrobiologie à la faculté des sciences de Meknès.

**Table1:** Characteristics of the different sampling sites

	Stations	Types of pollution	Lambert Contact Information		
			Latitude	Longitude	Altitude (meters)
<b>Bouishak watercourse</b>	B1	Domestic, agricultural and industrial	482120,6	364713,8	488
	B2	Domestic, agricultural and industrial	482205,4	365004,53	481
	B3	Domestic and industrial	482202,5	365900,2	458



**Figure 2:** Location of the study stations

The evaluation of the parasite load was determined by two techniques: modified Bailenger concentration technique strongly recommended by [15] and Faust technique (flootation method with zinc sulphate solution (33%, density,  $d = 1,18$ ) [16]. Egg identification has done using the WHO protocol (2004), the diagnostic manual for verminosis by coprology examination [17] and the manual of medical parasitology by Viviane GUILLAUME [18]. The criteria used for the identification of parasites are essentially: the size, shape, thickness of the shell, the presence or absence of polar plugs and a larva inside the egg. Quantification has done using a Mac Master slide under a light microscope. The total number per liter of Helminth eggs and protozoan cysts present in the wastewater sample is calculated using formula (1):

$$N = X / P \times V / S \quad (1)$$

With:  
 $N$  = number of eggs per liter of sample  
 $A$  = number of eggs counted on the McMaster blade or average number of eggs found in three blades  
 $X$  = volume of the final product (ml)  
 $P$  = McMaster Blade Capacity (0.3 ml)  
 $V$  = volume of the initial sample (1 liter).


### 3. Results and discussion

The results of parasitological analysis of wastewater from Bouishak wadi reveal a significant specific diversity of parasitic species. This is in agreement with other works [19-20].

This diversity made it possible to highlight the presence of thirteen parasite taxa (Table 2). Protozoan cysts (amoebae and intestinal flagellates) and Helminth eggs has divided into two classes: NEMATODES and CESTODES. The NEMATODES class is the most diversified with six taxa, a prevalence of 46% (Table 2) (Figure 3).The eggs of represent it: *Ascaris lumbricoides*, *Capillaria sp*, *Trichuris trichiura*, *Ancylostoma diudinal*, *Enterobius vermicularis*, and *Toxocara canis*. On the other hand, the class of CESTODES has a prevalence of 23% (3 taxa); the species collected are *Hymenolipis nana*, *Hymenolipis diminuta*, and *Taenia saginata*. The group of protozoa has represented by four taxa (31%): *Giardia lamblia*, *Entamoeba histolytica*, *Entamoeba coli* and *Blastocystis hominis*. It has noted that in all wastewater samples from different stations, protozoan cysts on helminth eggs are prominently predominant, consistent with previous work [20-21] (Figure 4). The number of Protozoan cysts is relatively high, ranging from  $6.49 \times 10^5$  cysts / L (station B3) to  $12.97 \times 10^5$  cysts / L (station B1), while station B2 contains  $9.62 \times 10^5$  cysts / L.

**Table2:** Inventory of the different species encountered in the urban wastewater of Bouishak collector.


Splitter	Classes	Species
Helminth	<i>Nematodes</i>	<i>Ascaris lumbricoides</i> <i>Trichuris trichiura</i> <i>Capillaria sp</i> <i>Toxocara canis</i> <i>Enterobius vermicularis</i> <i>Ancylostoma diudinal</i>
	<i>Cestodes</i>	<i>Hymenolipis nana</i> <i>Hymenolipis diminuta</i> <i>Taenia saginata</i>
	<i>Flagellated</i>	<i>Giardia lamblia</i> ,
	<i>Amoebas</i>	<i>Entamoeba coli</i> <i>Entamoeba histolytica</i> <i>Blastocystis hominis</i>
Protozoan		




**Figure 3:** Frequency of occurrence of parasite groups

The parasite load of the Helminth classes ranges from 1930 to 2651 eggs / L. The highest strength has recorded at station B1 while the lowest has obtained at station B3. Helminthes collected in Station B2 are in the order of 2314 eggs / L. On the other hand, the Helminths identified has characterized by the predominance of nematodes

with a frequency of 66% against 34% for the Cestodes (Figure 5). Our results are consistent with other work done in Morocco, Tunisia and France [21-22-23-24]. This unequal distribution of Nematodes with respect to Cestodes has explained by the fact that the first class is the most resistant in wastewater [21-25-26-27].




**Figure 4:** Distribution of parasites in the waters of Bouishak wadi




**Figure 5:** Frequency of occurrence of Helminth parasite classes

The high parasite load has detected at station B1. We has observed a decreasing slope from upstream (B1) to downstream (B3). This could explained by the fact that the B1 collector drains all domestic wastewater from a large agglomeration (Mansour, Camilia and Toulal) and the wastewater from the B2 collector is used by pumping for irrigation of the surrounding land, which has an impact on the parasitic load, noted low, at the B3 station (Figure 6). Within the Nematode class, *Ascaris lumbricoides* is the most dominant species in all stations (B1, B2 and B3) with an average concentration of about 114 eggs / L (Figure 6). Qualitatively, our results has corroborated with other works, this preponderance being due to the strong resistance of this kind to environmental conditions [28-29]. On the other hand, the presence of intestinal nematodes and particularly *Ascaris lumbricoides* and *Trichuris trichiura* in wastewater intended for irrigation constitutes a potential impact on human and animal health [28-36- 37-38]. Quantitative analysis of protozoan cysts in wastewater from Bouishak wadi revealed flagellates (*Giardia lamblia*) and amoebae (*Entamoeba coli*, *Entamoeba histolytica* and *Blastocystis hominis*) (Figure 7).




**Figure 6:** Egg concentration by class and taxon of wastewater from Bouishak wadi



**Figure 7:** Concentration of identified protozoa in wastewater from Bouishak wadi

We noted a predominance of *Giardia lamblia* cysts in the different stations and during the study period with average contents of  $39.6 \cdot 10^3$  cysts/L (B1),  $36.93 \cdot 10^3$  cysts/L (B2) and  $35.90 \cdot 10^3$  cysts/L (B3) (Figure 7). While *Entamoeba coli* cysts have the lowest concentration, from  $16.75 \cdot 10^3$  cysts/L (B1) to  $10.75 \cdot 10^3$  cysts/L (B3). From a quantitative point of view, the parasite load of the waters studied is large compared to that found by other researchers in other sites [21-30]. From a spatial point of view, mean cyst levels were high at station B1 with a total of  $10.8 \cdot 10^4$  cysts / L and increased to  $80.17 \cdot 10^3$  cysts / L then  $54.09 \cdot 10^3$  cysts/L respectively to B2 and B3 (Figure 8). This concentration, which decreases considerably from upstream to downstream, has explained by the decrease in the population size. The prevalence of *Giardia lamblia* and *Blastocystis hominis* in surface water and wastewater could lead to serious health risks, in this case giardiasis, which has manifested by watery diarrhea, steatorrhea and malabsorption, is a common infection in children [34].



**Figure 8:** Spatial variation of protozoan cyst densities

For example, the presence of intestinal nematodes, particularly *Ascaris lumbricoides*, and *Trichuris trichiura* in wastewater intended for irrigation has considered a major health and environmental hazard [35-36-37]. The World Health Organization (WHO) [38] estimated the number of cases of Ascaridiosis due to *Ascaris lumbricoides* and 500 million cases of *Trichuris trichiura* infection in the world at 1,000 million. In 1994, Chan et al., [39] estimated that there were 1471 million and 1048 million cases of *A. lumbricoides* and *T. trichiura* infections in the world, respectively. Clinically, *A. lumbricoides* can cause bowel obstruction and *T. trichiura* has been associated with dysentery [40]. Both Helminths have also been associated with stunting [41-42-43]. Massive infestations of these two nematodes are more common in children than in adults.

## Conclusion

The results of our study show that helminth protozoan egg and cyst concentrations far exceed the standards recommended by the WHO guidelines and Moroccan water standards for irrigation described in the decree. Ministerial Meeting No. 1276-01 of the year 2002. Indeed, it claims a parasite load less than or equal to a viable egg per liter for water for irrigation crops. A qualitative and quantitative divergence of sewage pests has noted. Sewage upstream of Bouishak wadi was more parasitic than downstream water. This difference in concentration may be due to the size and socio-economic level of the urban population, the geographical context, the arid and semi-arid climate of the region, and the urban effluents that has discharged daily into the river. wadi. Therefore, the control of waterborne diseases becomes an essential strategy to reduce risk factors.

## References

1. M. Bentaleb, Réutilisation des eaux usées brutes en irrigation au Maroc. Mémoire de troisième cycle, Option Génie Rural, Institut Agronomique et Vétérinaire, Hassan II, Maroc Institut Agronomique et Vétérinaire Hassan II, Rabat (1986).
2. H. Abouzeid, L'état de l'expérimentation et de la recherche en matière de réutilisation des eaux usées au Maroc. *Rev. Eau & Development* (10) (1990) 6-11.
3. A. Ait melloul, O. Amahmid, L. Hassani, K. Bouhoum. Health effect of human wastes use in agriculture in El Azzouzia (the wastewater spreading area of Marrakech city, Morocco). *Inter. J. Envir Health Research* 12 (2002) 17– 23.
4. H. Talouizte, M. Merzouki, A. El Ouali Lalami, L. Bennani, M. Benlemlih, Evolution de la charge microbienne de la laitue irriguée avec les eaux usées urbaines de la ville de Fès au Maroc, *Tribune de l'eau*. 642 (2007)
5. OMS (Organisation Mondiale de la Santé), *Recommendations*, 3rd, Ed. World Health Organization. Geneva Vol.1 (2004).

6. A. Aboulkacem. . Etude comparative de la qualité bactériologique des eaux des oueds Boufekrane et Ouislane à la traversée de la ville de Meknès (Maroc). *Revue Microbiol. Ind.San.et Environ* 1 (2007) 10-22.
- 7 J. Addouli, A. Chahlaoui, A. Berrahou, A. Chafi, A. Ennabili, L. Karrouch, Influence des eaux usées, utilisées en irrigation, sur la qualite des eaux de l'Oeud Bouishak - Region de Meknes (Centre-Sud du Maroc) *Revue Microbiol. Ind. Santé et Environn.* 3, N°1 (2009) 56-75.
8. A. Chahlaoui, Vulnérabilité des nappes superficielles et subaffleurantes du sud-ouest tunisien. *Sécheresse*, 2 N°11 (2000) 85-91.
9. RADEEM (Régie Autonome de Distribution de l'Eau et de l'Electricité), (2003).
10. M. Abdouh, M. El Atrouz, A. Mechkouri. Profil environnemental de Meknès. (2004)28-29.
11. RGPH (Recensement Général de la Population et de l'Habitat) (2014)
12. M. Cadillon, Y. Reaumaux, J. Bize, Réutilisation des eaux usées : Contraintes et enjeux. Actes des journées techniques d'assainissement au Maroc, Agadir (1989) 2-26.
13. R. Firadi, Epuration et réutilisation des eaux usées de la ville d'Ouarzazate en agriculture : devenir des œufs d'helminthes et étude de leur viabilité dans les eaux usées et les boues. Thèse de 3èmecycle en Biologie, Option : Traitement et valorisation agricole des eaux usées, Université Cadi Ayyad –Faculté des Sciences Semlalia, Marrakech, (1996)114p.
14. K. Ouarrak, A. Chahlaoui and I. Taha, Monitoring of the parasitic load wastewater of Ouislane wadi and Bouishak wadi from the city of meknes moroccoijas, *Inter.J. of Innovation Sciences and Research*. 6 N°7(2017) 1063-1067
15. OMS (Organisation mondiale de la Santé). Planches pour le diagnostic des parasites intestinaux. Genève, (1994) 29 p.
16. J. Rodier, C. Bazin, J.P. Broutin, P. Chambon, H. Champsaur, L. Rodi, l'analyse de l'eau naturelle, eau résiduaires, eau de mer 8e édition. *Duno, Paris, France* (2009).
17. D. Thienpont, M. Cadillon, Y. Reaumaux, and J. Bize, Le traitement et valorisation des eaux usées d'Oujda. Rapport préliminaire du programme de coopération Franco-marocaine, collectivités locales (1993).
18. Gu. Viviane, Parasitologie, De boeck, Paris. *Livre* (2007) 182p
19. C. Aliaume, G. Lasserre, M. Louis, Organisation spatial des peuplements ichtyologiques des herbiers à Thalassia du Grand Culde-sac marin en Guadeloupe. *Revue d'hydrobiologie Tropicale*, 23 : 231- 250. *Revue d'hydrobiologie Tropicale*, 23 (1990) 231- 250.
20. R. Stott, T. Jenkins, M. Shabana & E. May, A survey of the microbial quality of wastewater in Ismailia, Egypt and the implications for wastewater reuse. *Sc. Tech.* 35(1997) 211-217.
21. I. Sylla, D. Belghyti, Analyse parasitologique des eaux brutes de la ville de Sidi Yahia Gharb (Maroc). Revue Mondiale de la Recherche Biologique, *World Journal of Biological Research*, ISSN : 10 (2008) 1994-5108 ISPROMS,
22. Z. Alouini, Flux de la charge parasitaire dans cinq stations d'épuration en Tunisie. *Revue des Sciences de l'eau* 6 N°4 (1993) 453-62. <https://doi.org/10.7202/705185ar>.
23. Z. Alouini, H.Achour, A. Alouini, Devenir de la charge parasitaire des eaux usées traitées dans le réseau d'irrigation "Cebala" in Zekri, Laajini A, agriculture durabilité et environnement. CIHEAM (1995)117-124.
24. J.L. Stien, J. Schwartzbrod, Devenir des œufs d'helminthes au cours d'un cycle d'épuration des eaux usées urbaines, *Revue internationale des séries de l'eau*. T.S.M., N°IO, pp 495-498. *Revue internationale des séries de l'eau*, 3 (3/4) (1987) 77-82.
25. M. Guessab, J. Bize, J. Schwartzbrod, A. Mani, M. Morlot, N. Nivault, L. Schwartzbrod, Wasterwater treatment dy infiltration percolation on sand: results in Ben Sergao. Morocco, *Sci. Tech.* 17(1993)91-95
26. J. Schwartzbrod, S. Banas, Parasite contamination of liquid sludge from urban wastewater treatment plants. Water, *Sci. Tech.* Vol. 47 N°3. (2003)163-166
27. K. Hajjami, M. M. Ennaji, S. Fouad, N. Oubrim, K. Khallayoune, N. Cohen. Wastewater Reuse for Irrigation in Morocco: Helminth Eggs Contamination's Level of Irrigated Crops and Sanitary Risk (A Case Study of Settat and Soualem Regions) *Journal of Bacteriology and Parasitology*. 2 N° 5 (2012)193-201.
28. Y. El Guamri, D. Belghyti, Etude de la qualité physico-chimique des eaux usées brutes de la commune urbaine de Saknia, rejetées dans le lac Fouarat (Kénitra, Maroc) : *Journal Africain des Sciences de l'Environnement* 1 (2006) 53-60 *Afrique SCIENCE* 03(1) (2007)123 – 145.
29. S. Raweh, K. Elkharrim, M. Cissé, Y. El Guamri, Y. Abchir, D. Belghyti, Aspect parasitologique des eaux usées rejetées dans le bas Sebou (Kenitra, Maroc). *World Journal of Biological Research*. 3 (1) (2010) 18-29.

- 30 N. Rhallabi, R. Moundib, M. Maaroufy, M. Marghich, Kh. Khalaayoune, Kh. Bouzoubaa, A. Agoumi, M. Ettalibi, H. Berdai, N. Jellal, M. Touzani, M. Mekrane, A. Handouf, T. Benchokroum, B. EL Hamouri,. *Actes Inst. Agron. Vet.* 10 (1990)57-66.
31. L.W. Casson, C.A. Sorber, J.L. Sykora, P.D. Gavaghan, M.A. Shapiro, W. Jakubowski, Giardia in Wastewater-Effect of Treatment Research *Journal of the Water Pollution Control Federation* 62 N°5 (1990) 670-675.
32. K. Bouhoum, O. Amahmid, S. Asmama. Devenir des œufs d'helminthes et des kystes de protozoaires dans un canal à ciel ouvert alimenté par les eaux usées de Marrakech. *Revue des Sciences de l'Eau*, 10 N°2 (1997) 147-246 ; <https://doi.org/10.7202/705278ar>
33. M. Abouelouafa, H. El Halouani, M. Kharboua et A. Berrichi. 2002. Caractérisation physicochimique et bactériologique des eaux usées brutes de la ville d'Oujda : Canal principal et Oued Bounâîm. *Actes Inst. Agron. Vet. (Maroc)* Vol. 22 (3) (2002) 143 - 150.
34. Wright SG. Protozoan infections of the gastrointestinal tract. *Infect Dis Clin North Am.* (2012) 26(2):323-39. <https://doi.org/10.1016/j.idc.2012.03.009>.
35. M.I. Shuval, A. Adin, B. Fatal, E. Rawitz, P. Yekutiel. Wastewater irrigation in developing countries: Health effects and technical solutions' *World Bank Technical*, paper N° 51 (1986) 5121-38.
36. D. Blum, R.G. Feachem. 1985. Health aspects of night soil and sludge use in agriculture. Part III: an epidemiological perspective. Report N° 05/85 (1985). *Part III. Dübendorf, international référence centre for waste disposai. Report N° 05/85*
37. M. Straus, Survival of excreted pathogens in excreta and faecal sludges. *IRCWD NEWS* 23(1985) 4-9.
38. Organisation Mondiale de la Santé (OMS). L'utilisation des eaux usées en agriculture et en aquaculture : recommandations à avisées sanitaires. Rapport d'un groupe scientifique de l'OMS. Organisation mondiale de la santé, série de rapports techniques, n° 778, Genève (1989).
39. L. Chan, DA, Bundy, SP. Kan. Nematode Infections Are Risk Factors for Staphylococcal Infection in Children Vol. 97 (3) (2002) 395-399, April; <http://dx.doi.org/10.1590/S0074-02762002000300021>
40. E.S. Cooper Whyte-Alleng, C.A.M, JS,Finzi-Smith et T.T. MacDonald, Intestinal nematode infections in children: the pathophysiological price paid. *Parasitology* 104 (Suppl S1) (1992) S91-S103. <https://doi.org/10.1017/S0031182000075272>
41. D. A. P. BUNDY, E. S. COOPER, D. E. THOMPSON, J. M. DIDIER, Effect of age and initial infection status on the rate of reinfection with *Trichuris trichiura* after treatment. *Parasitology* 97 N°3 (1988) 469–476. <https://doi.org/10.1017/S003118200005887X>
42. LS. Stephenson, MC. Latham, EJ. Adams, SN, Kinoti , A. Pertet. *J Nutr* Physical fitness, growth and appetite of Kenyan school boy's .ith hookworm, Trichuris trichiura and Ascaris lumbricoides infections are improved four months after a single dose of albendazole. 123(6) (1993) 1036-46 <https://doi.org/10.1093/jn/123.6.1036>
43. JW. Lawless , Latham MC, Stephenson LS, Kinoti SN, Pertet AM, Iron supplementation improves appetite and growth in anemic Kenyan primary school children 124 N° 5 (1994) 645-654 <https://doi.org/10.1093/jn/124.5.645>.

(2019) ; <http://www.jmaterenvironsci.com>